Jacobs University Bremen

Thesis for Bachelor of Science in Mathematics

by Mengyuan Zhang

arXiv:1406.4753v1 [math.RT] 18 Jun 2014

Lie algebras of linear systems and their automorphisms

Advisor
Prof. Dr. Ivan Penkov

Abstract

The objective of this thesis is to study the automorphism groups of the Lie algebras attached to linear systems. A linear system is a pair of vector spaces (U, W) with a nondegenerate pairing $\langle \cdot, \cdot \rangle \colon U \otimes W \to \mathbb{C}$, to which we attach three Lie algebras $\mathfrak{sl}_{U,W} \subset \mathfrak{gl}_{U,W} \subset \mathfrak{gl}^M_{U,W}$. If both U and W are countable dimensional, then, up to isomorphism, there is a unique linear system (V, V_*). In this case $\mathfrak{sl}_{V,V_*}$ and $\mathfrak{gl}_{V,V_*}$ are the well-known Lie algebras $\mathfrak{sl}_\infty$ and $\mathfrak{gl}_\infty$, while the Lie algebra $\mathfrak{gl}^M_{V,V_*}$ is the Mackey Lie algebra introduced in [PS13].

We review results about the monoidal categories $\mathbb{T}_{\mathfrak{sl}_{U,W}}$ and $\mathbb{T}_{\mathfrak{gl}^M_{U,W}}$ of tensor modules, both of which turn out to be equivalent as monoidal categories to the category $\mathbb{T}_{\mathfrak{sl}_\infty}$ introduced earlier in [DCPS11]. Using the relations between the categories $\mathbb{T}_{\mathfrak{sl}_\infty}$ and $\mathbb{T}_{\mathfrak{gl}^M_\infty}$, we compute the automorphism group of $\mathfrak{gl}^M_\infty$.

Contents

1 Preliminaries **2**

2 The Lie algebras $\mathfrak{sl}_{U,W}$ and $\mathfrak{gl}_{U,W}$ **3**

3 The Mackey Lie algebras $\mathfrak{gl}^M_{U,W}$ **6**

4 The categories $\mathbb{T}_{\mathfrak{sl}_{U,W}}$, $\mathbb{T}_{\mathfrak{gl}^M_{U,W}}$ and the functor $\mathcal{F}^h$ **8**

5 Automorphisms of $\mathfrak{sl}_\infty$ and $\mathfrak{gl}^M_\infty$ **11**

1 Preliminaries

The ground field is $\mathbb{C}$. $(\,\cdot\,)^*$ denotes the contravariant functor which maps a vector space M to $\mathrm{Hom}(M,\mathbb{C})$ and maps a linear map $f\colon M \to N$ to its dual $f^*\colon N^* \to M^*$. In this paper, countable dimensional means infinite countable dimensional. $T(X)$ denotes the tensor algebra of a vector space X. $\mathfrak{g}$ denotes an arbitrary Lie algebra, and $\mathfrak{g}$-mod denotes the category of $\mathfrak{g}$-modules, where morphisms are $\mathfrak{g}$-homomorphisms.

Throughout this paper, let V be a fixed vector space with a fixed countable basis $\mathcal{B} = \{e_i\}_{i\in\mathbb{N}}$ and let e^i be the linear functional dual to e_i, i.e. $e^i(e_j) = \delta_{ij} \quad \forall i,j$. Let $\mathcal{B}^* := \{e^i\}_{i\in\mathbb{N}}$ and $V_* := \operatorname{span}\mathcal{B}^*$. We also set $\mathcal{B}_n := \{e_i\}_{i=1}^n$ and $\mathcal{B}_n^* := \{e^i\}_{i=1}^n$. Then clearly $V = \varinjlim V_n$ and $V_* = \varinjlim V_n^*$, where $V_n := \operatorname{span}\mathcal{B}_n$ and $V_n^* := \operatorname{span}\mathcal{B}_n^*$.

Mat_n denotes the Lie algebra of $n\times n$ matrices. For $n \le m$, consider the embedding $\mathrm{Mat}_n \xhookrightarrow{\iota_{nm}} \mathrm{Mat}_m$ by upper left corner identity inclusion and filling zeros elsewhere. Then $(\{\mathrm{Mat}_n\},\{\iota_{nm}\})$ forms a direct system of Lie algebras, and $\mathrm{Mat}_\mathbb{N} := \varinjlim \mathrm{Mat}_n$ is its direct limit. $\mathrm{Mat}_\mathbb{N}$ is nothing but the Lie algebra of matrices $(A_{ij})_{i,j\in\mathbb{N}}$ that have finitely many nonzero entries.

We define the Lie algebra $\mathfrak{gl}_n := V_n \otimes V_n^*$, with Lie bracket given by

$$[e_i\otimes e^j, e_k\otimes e^l] = \delta_{jk}e_i\otimes e^l - \delta_{il}e_k\otimes e^j \quad \forall e_i,e_k\in\mathcal{B}_n, e^j,e^l\in\mathcal{B}_n^*. \tag{1}$$

Observe that $\mathfrak{gl}_n$ and Mat_n are isomorphic as Lie algebras. An isomorphism is given by sending $e_i\otimes e^j$ to the matrix E_{ij} having 1 at (i,j) and 0 elsewhere. For any $n\le m$, the embedding ι_{nm} induces an imbedding $\mathfrak{gl}_n \xhookrightarrow{\iota'_{nm}} \mathfrak{gl}_m$ such that the following diagram commutes

$$\begin{array}{ccc} \mathfrak{gl}_n & \xhookrightarrow{\iota'_{nm}} & \mathfrak{gl}_m \\ \updownarrow & & \updownarrow \\ \mathrm{Mat}_n & \xhookrightarrow{\iota_{nm}} & \mathrm{Mat}_m, \end{array}$$

making $(\{\mathfrak{gl}_n\},\{\iota'_{nm}\})$ a direct system with direct limit

$$\mathfrak{gl}_\infty := V\otimes V_* = \varinjlim(V_n\otimes V_n^*) = \varinjlim \mathfrak{gl}_n.$$

The Lie bracket on $\mathfrak{gl}_\infty$ is given by (1), where now $e_i,e_k\in\mathcal{B}, e^j,e^l\in\mathcal{B}^*$.

Clearly, since the two direct systems $(\{\mathrm{Mat}_n\},\{\iota_{nm}\})$ and $(\{\mathfrak{gl}_n\},\{\iota'_{nm}\})$ are isomorphic, their direct limits $\mathrm{Mat}_\mathbb{N}$ and $\mathfrak{gl}_\infty$ are isomorphic.

The trace homomorphism $\mathrm{tr}_n\colon \mathfrak{gl}_n \to \mathbb{C}$ can be extended to a homomorphism $\mathrm{tr}\colon \mathfrak{gl}_\infty \to \mathbb{C}$. Define $\mathfrak{sl}_\infty := \ker \mathrm{tr}$. Then $\mathfrak{sl}_\infty = \varinjlim \ker \mathrm{tr}_n$. Similarly, $\mathfrak{sl}_n \cong \mathrm{Mat}_n^0$ and $\mathfrak{sl}_\infty \cong \mathrm{Mat}_\mathbb{N}^0$, where Mat_n^0 denotes the Lie subalgebra of Mat_n consisting of traceless $n\times n$ matrices and $\mathrm{Mat}_\mathbb{N}^0$ denotes the Lie subalgebra of $\mathrm{Mat}_\mathbb{N}$ consisting of traceless finitary matrices.

2 The Lie algebras $\mathfrak{sl}_{U,W}$ and $\mathfrak{gl}_{U,W}$

In this section we generalize the construction of $\mathfrak{sl}_\infty$ and $\mathfrak{gl}_\infty$ to arbitrary linear systems (U, W). A pair of vector spaces of arbitrary dimensions (U, W) is called a *linear system* if they are equipped with a nondegenerate bilinear form $\langle \cdot, \cdot \rangle \colon U \times W \to \mathbb{C}$. We say that (U', W') is a *subsystem* if U', W' are subspaces of U, W respectively and $\langle \cdot, \cdot \rangle$ is nondegenerate when restricted to $U' \times W'$.

Proposition 2.1. Let (U, W) be a linear system and let $U_f \subset U$ be a finite-dimensional subspace of U. Let W_f be a direct complement to $U_f^\perp$ in W, i.e. $W = U_f^\perp \oplus W_f$. Then (U_f, W_f) is a subsystem and $\dim U_f = \dim W_f$.

Proof. Suppose $v \in V_f$ is such that $\langle v, W_f \rangle = 0$. Then $\langle v, W \rangle = 0$, and we must have $v = 0$. Suppose $w \in W_f$ is such that $\langle V_f, w \rangle = 0$. Then $w \in V_f^\perp \cap W_f$, thus $w = 0$. We conclude that $\langle \cdot, \cdot \rangle$ is nondegenerate on $V_f \times W_f$. Therefore the nondegenerate form $\langle \cdot, \cdot \rangle$ induces an injection $W_f \hookrightarrow (V_f)^*$. This forces W_f to be finite-dimensional. Similarly, there is an injection $V_f \hookrightarrow (W_f)^*$. Therefore $\dim V_f = \dim W_f$. □

Given any linear system (U, W), we define two Lie algebras $\mathfrak{gl}_{U,W}$ and $\mathfrak{sl}_{U,W}$. Let $\mathfrak{gl}_{U,W}$ equal the vector space $U \otimes W$, with Lie bracket

$$[u_1 \otimes w_1, u_2 \otimes w_2] = \langle u_2, w_1 \rangle u_1 \otimes w_2 - \langle u_1, w_2 \rangle u_2 \otimes w_1 \quad \forall u_1, u_2 \in U, w_1, w_2 \in W.$$

We define $\mathfrak{sl}_{U,W}$ to be the kernel of $\langle \cdot, \cdot \rangle$. Note that the pair (V, V_*) is a linear system, and $\mathfrak{sl}_{V,V_*} = \mathfrak{sl}_\infty$ and $\mathfrak{gl}_{V,V_*} = \mathfrak{gl}_\infty$. The next four propositions generalize some simple observations concerning $\mathfrak{sl}_\infty$ and $\mathfrak{gl}_\infty$ to arbitrary $\mathfrak{sl}_{U,W}$ and $\mathfrak{gl}_{U,W}$.

Proposition 2.2. There are isomorphisms of Lie algebras $\mathfrak{sl}_{U,W} \cong \mathfrak{sl}_{W,U}$ and $\mathfrak{gl}_{U,W} \cong \mathfrak{gl}_{W,U}$.

Proof. Consider the linear operators $f \colon \mathfrak{gl}_{U,W} \to \mathfrak{gl}_{W,U}$ and $g \colon \mathfrak{gl}_{W,U} \to \mathfrak{gl}_{U,W}$ such that

$$f(u \otimes w) = -w \otimes u \quad \forall u \otimes w \in \mathfrak{gl}_{U,W},$$

$$g(w \otimes u) = -u \otimes w \quad \forall u \otimes w \in \mathfrak{gl}_{W,U}.$$

They are both Lie algebra homomorphisms and are mutually inverse. Also f restricts to an isomorphism of Lie algebras $\mathfrak{sl}_{U,W} \cong \mathfrak{sl}_{W,U}$. □

Proposition 2.3. The Lie algebra $\mathfrak{sl}_{U,W}$ is simple.

Proof. The set of finite-dimensional subsystems (U_f, W_f) of (U, W) is partially ordered by inclusion, and any two such subsystems have an upper bound. Thus we obtain a direct system of Lie algebras $\{\mathfrak{sl}_{U_f,W_f}\}$ with direct limit $\mathfrak{sl}_{U,W} = \varinjlim \mathfrak{sl}_{U_f,W_f}$. Any nontrivial ideal I of $\mathfrak{sl}_{U,W}$ intersects nontrivially with some $\mathfrak{sl}_{U_f,W_f}$. We conclude that $I \supset \mathfrak{sl}_{U_f,W_f}$ by the simplicity of $\mathfrak{sl}_{U_f,W_f}$. Similarly, $I \supset \mathfrak{sl}_{U_f',W_f'}$ for any finite-dimensional subsystem (U_f', W_f') containing (U_f, W_f) by the simplicity of $\mathfrak{sl}_{U_f',W_f'}$. This means that $I = \mathfrak{sl}_{U,W}$. □

Proposition 2.4. The Lie algebra $\mathfrak{sl}_{U,W}$ is the commutator subalgebra of $\mathfrak{gl}_{U,W}$.

Proof. It is clear from the definition of the Lie bracket that $[\mathfrak{gl}_{U,W}, \mathfrak{gl}_{U,W}] \subset \mathfrak{sl}_{U,W}$. Conversely, observe that $[\mathfrak{sl}_{U,W}, \mathfrak{sl}_{U,W}]$ is a nontrivial ideal of $\mathfrak{sl}_{U,W}$, therefore equal to $\mathfrak{sl}_{U,W}$ by the simplicity of $\mathfrak{sl}_{U,W}$. To conclude,

$$\mathfrak{sl}_{U,W} = [\mathfrak{sl}_{U,W}, \mathfrak{sl}_{U,W}] = [\mathfrak{gl}_{U,W}, \mathfrak{gl}_{U,W}].$$ □

Proposition 2.5.

1. Let A be any finite-dimensional Lie subalgebra of $\mathfrak{gl}_{U,W}$, then $A \subset \mathfrak{gl}_{U_f,W_f}$ for some finite-dimensional subsystem (U_f, W_f).
2. Let A be any finite-dimensional Lie subalgebra of $\mathfrak{sl}_{U,W}$, then $A \subset \mathfrak{sl}_{U_f,W_f}$ for some finite-dimensional subsystem (U_f, W_f).

Proof.

1. Let $\{u_\alpha\}$ be a basis of U and $\{w_\beta\}$ be a basis of W. Then $\{u_\alpha \otimes w_\beta\}$ is a basis of $U \otimes W$. Fix a basis $\{a_1, \cdots, a_n\}$ of A, and let

$$a_i = \sum_{k=1}^{n_i} c_{i,k} \cdot u_{\alpha_{i,k}} \otimes w_{\beta_{i,k}}.$$

   Let (U_f, W_f) be a finite-dimensional subsystem such that $U_f \supset \mathrm{Span}\{u_{\alpha_{i,k}}\}$ and $W_f \supset \mathrm{Span}\{w_{\beta_{i,k}}\}$. Then $A \subset \mathrm{Span}\{u_{\alpha_{i,k}}\} \otimes \mathrm{Span}\{w_{\beta_{i,k}}\} \subset \mathfrak{gl}_{U_f,W_f}$.
2. Given a finite-dimensional subalgebra $A \subset \mathfrak{sl}_{U,W} \subset \mathfrak{gl}_{U,W}$, we have $A \subset \mathfrak{gl}_{U_f,W_f}$ for some finite-dimensional subsystem (U_f, W_f) by statement 1. Then $A \subset \mathfrak{gl}_{U_f,W_f} \cap \ker\langle\cdot,\cdot\rangle = \mathfrak{sl}_{U_f,W_f}$. □

Having defined $\mathfrak{sl}_{U,W}$ and $\mathfrak{gl}_{U,W}$ for a linear system (U, W), it is natural to ask when are $\mathfrak{gl}_{U_1,W_1}$ and $\mathfrak{gl}_{U_2,W_2}$ isomorphic as Lie algebras. A necessary and sufficient condition is given in [PS13].

Definition. Two linear systems (U_1, W_1) and (U_2, W_2) are isomorphic iff one of the following holds:

1. There are vector space isomorphisms $f\colon U_1 \to U_2$ and $g\colon W_1 \to W_2$ such that $\langle f(u), g(w)\rangle = \langle u, w\rangle$ for all $u \in U_1, w \in W_1$.
2. There are vector space isomorphisms $f\colon U_1 \to W_2$ and $g\colon W_1 \to U_2$ such that $\langle g(w), f(u)\rangle = \langle u, w\rangle$ for all $u \in U_1, w \in W_1$.

We write $(U_1, W_1) \cong (U_2, W_2)$.

Theorem 2.6. [PS13, Prop 1.1] The Lie algebras $\mathfrak{sl}_{(U_1,W_1)}$ and $\mathfrak{sl}_{(U_2,W_2)}$ are isomorphic iff the linear systems (U_1, W_1) and (U_2, W_2) are isomorphic.

Corollary 2.7. The Lie algebras $\mathfrak{gl}_{(U_1,W_1)}$ and $\mathfrak{gl}_{(U_2,W_2)}$ are isomorphic iff the linear systems (U_1, W_1) and (U_2, W_2) are isomorphic.

Proof. $\Longleftarrow$ If we have a linear system isomorphism $f\colon U_1 \to U_2$ and $g\colon W_1 \to W_2$, then

$$f \otimes g\colon u \otimes w \mapsto f(u) \otimes g(w) \quad \forall u \otimes w \in \mathfrak{gl}_{U_1,W_1}$$

induces an isomorphism of Lie algebras. If we have isomorphisms $f\colon U_1 \to W_2$ and $g\colon W_1 \to U_2$, then

$$-g \otimes f\colon u \otimes w \mapsto -g(w) \otimes f(u) \quad \forall u \otimes w \in \mathfrak{gl}_{U_1,W_1}$$

induces an isomorphism of Lie algebras.

$\Longrightarrow$ Conversely, if $\mathfrak{gl}_{U_1,W_1} \cong \mathfrak{gl}_{U_2,W_2}$ then the commutator subalgebras $\mathfrak{sl}_{U_1,W_1}$ and $\mathfrak{sl}_{U_2,W_2}$ are isomorphic. By Proposition 2.6 we conclude that $(U_1, W_1) \cong (U_2, W_2)$. □

We call a linear system (U, W) *countable dimensional* if both U and W are countable dimensional. In such cases we can say more about $\mathfrak{gl}_{U,W}$ and $\mathfrak{sl}_{U,W}$ because of the following observation in [Mac43]. The next theorem and corollary tell us that, up to isomorphism, there is only one countable dimensional linear system.

Theorem 2.8. [Mac43] For any countable dimensional linear system (U, W), there exist dual bases $\{\tilde{u}_i\}_{i\in\mathbb{N}}$ of U and $\{\tilde{w}_i\}_{i\in\mathbb{N}}$ of W, where $\langle \tilde{u}_i, \tilde{w}_j\rangle = \delta_{ij}\ \forall i, j$.

Proof. Start with any basis $\{u_i\}_{i\in\mathbb{N}}$ of U and $\{w_i\}_{i\in\mathbb{N}}$ of W and perform the Gram-Schmidt algorithm.

Step 1: If $\langle u_1, w_1\rangle \neq 0$ then go to step 2. If not, then by non-degeneracy of the form $\langle\cdot,\cdot\rangle$ there is a w_j such that $\langle u_1, w_j\rangle \neq 0$. Replace w_1 by $w_1 + w_j$.

Step 2: Set $\tilde{u}_1 = u_1$. Adjust w_1 by a scalar to obtain $\tilde{w}_1$ such that $\langle \tilde{u}_1, \tilde{w}_1\rangle = 1$. Replace u_2 by $\tilde{u}_2 = u_2 - \langle u_2, \tilde{w}_1\rangle\tilde{u}_1$, whence $\langle \tilde{u}_2, \tilde{w}_1\rangle = 0$. Now adjust w_2 to make sure $\langle \tilde{u}_2, w_2\rangle = 1$ and set $\tilde{w}_2 = w_2 - \langle u_1, w_2\rangle\tilde{w}_1$. We have $\langle \tilde{u}_1, \tilde{w}_2\rangle = 0$ and $\langle \tilde{u}_2, \tilde{w}_2\rangle = \langle \tilde{u}_2, w_2\rangle - \langle u_1, w_2\rangle\langle \tilde{u}_2, \tilde{w}_1\rangle = 1$.

Step 3: Having obtained $\tilde{u}_1, \cdots, \tilde{u}_n$ and $\tilde{w}_1, \cdots, \tilde{w}_n$, we set

$$\tilde{u}_{n+1} = u_{n+1} - \sum_{i=1}^{n} \langle u_{n+1}, \tilde{w}_i\rangle\tilde{u}_i,$$

and apply argument in step 1 to adjust w_{n+1} such that $\langle \tilde{u}_{n+1}, w_{n+1}\rangle = 1$. We also set

$$\tilde{w}_{n+1} = w_{n+1} - \sum_{i=1}^{n} \langle \tilde{u}_i, w_{n+1}\rangle\tilde{w}_i.$$

By induction this will give us the dual bases. □

Corollary 2.9. If (U, W) is countable dimensional, then $(U, W) \cong (V, V_*)$. As a consequence $\mathfrak{sl}_{U,W} \cong \mathfrak{sl}_\infty$ and $\mathfrak{gl}_{U,W} \cong \mathfrak{gl}_\infty$.

Proof. Given a pair of dual bases $\{u_n\}_{n\in\mathbb{N}}$ of U and $\{w_n\}_{n\in\mathbb{N}}$ of W, the isomorphisms $f : U \to V$ and $g : W \to V_*$ defined by $f(u_n) = e_n$ and $g(w_n) = e^n$ preserve the form $\langle\cdot,\cdot\rangle$. Therefore the linear systems (U, W) and (V, V_*) are isomorphic. By Theorem 2.6 and Corollary 2.7, we conclude that there are isomorphisms of Lie algebras $\mathfrak{sl}_{U,W} \cong \mathfrak{sl}_\infty$ and $\mathfrak{gl}_{U,W} \cong \mathfrak{gl}_\infty$. □

Theorem 2.8 clearly holds for finite-dimensional linear systems as well, but fails for uncountable dimensional linear systems. Consider $(X := W \oplus W^*, X' := W \oplus W^*)$ where W is a countable dimensional vector space and the form $X \times X' \to \mathbb{C}$ is given by

$$(w + w^*, v + v^*) \mapsto w^*(v) + v^*(w) \quad \forall v, w \in W, v^*, w^* \in W^*.$$

If there were a pair of dual bases, then $W^\perp \subset X'$ would be uncountable dimensional. However, we observe that $W^\perp = W \subset X'$.

3 The Mackey Lie algebras $\mathfrak{gl}^M_{U,W}$

In this section we introduce a canonical Lie algebra $\mathfrak{gl}^M_{U,W}$ corresponding to the linear system (U, W). We discuss the relations between $\mathfrak{sl}_{U,W}, \mathfrak{gl}_{U,W}$ and $\mathfrak{gl}^M_{U,W}$, and realize U and W as non-isomorphic modules of $\mathfrak{gl}^M_{U,W}$.

Definition. We embed $W \hookrightarrow U^*$ and $U \hookrightarrow W^*$ using the pairing $\langle \cdot, \cdot \rangle$. Set

$$\mathfrak{gl}^M_{U,W} := \{\varphi \in \mathrm{End}(U) |\ \varphi^*(W) \subset W\}, \quad \mathfrak{gl}^M_{W,U} := \{\psi \in \mathrm{End}(W) |\ \psi^*(U) \subset U\}.$$

Then $\mathfrak{gl}^M_{U,W}$ and $\mathfrak{gl}^M_{W,U}$ are Lie subalgebras of the Lie algebras $\mathrm{End}(U)$ and $\mathrm{End}(W)$, respectively.

It's easy to see that there is an isomorphism $\mathfrak{gl}^M_{U,W} \cong \mathfrak{gl}^M_{W,U}$ of Lie algebras. The map that sends $\varphi \in \mathfrak{gl}^M_{U,W}$ to $-\varphi^*_{|W}$, is a Lie algebra homomorphism whose inverse homomorphism is given by sending $\psi \in \mathfrak{gl}^M_{W,U}$ to $-\psi^*_{|U}$.

Proposition 3.1 (Penkov, Serganova)**.** The Lie algebra $\mathfrak{gl}_{U,W}$ is isomorphic to the ideal S of $\mathfrak{gl}^M_{U,W}$, where

$$S := \{\varphi \in \mathfrak{gl}^M_{U,W} \colon \dim \varphi(U) < \infty, \dim \varphi^*(W) < \infty\}.$$

Proof. First of all, it is clear that $[h, \varphi] = h\varphi - \varphi h$ has finite-dimensional image in both U and W if one of h, φ does. Thus, S is an ideal. Consider the injection of Lie algebras $\iota \colon \mathfrak{gl}_{U,W} \to \mathfrak{gl}^M_{U,W}$ induced by $u \otimes w \mapsto \langle \cdot, w \rangle u$. The range of ι is indeed in $\mathfrak{gl}^M_{U,W}$ because

$$(\iota(u \otimes w))^*(\langle \cdot, y \rangle) = \langle \langle \cdot, w \rangle u, y \rangle = \langle \cdot, \langle u, y \rangle w \rangle \in W \quad \forall \langle \cdot, y \rangle \in W.$$

Moreover, since elements of $U \otimes W$ are finite linear combinations of pure tensors, under ι they have finite-dimensional images in both U and W, which means that $\mathrm{im}\, \iota \subset S$.

We now show that $\mathrm{im}\, \iota \supset S$. Let (U_f, W_f) be a finite-dimensional subsystem of (U, W). Consider the Lie subalgebra

$$S_{U_f, W_f} := \{\varphi \in \mathrm{End}(U) \colon \varphi(U) \subset U_f \text{ and } \varphi^*(W) \subset W_f\} \subset \mathfrak{gl}^M_{U,W}.$$

Observe that $\langle \varphi(v), W \rangle = \langle v, \varphi^*(W) \rangle = 0$ for all $v \in W_f^\perp$. Therefore $U \supset \ker \varphi \supset W_f^\perp$ for any $\varphi \in S_{U_f, W_f}$. In addition, since $\varphi(U) \subset U_f$, the subspace S_{U_f, W_f} can at most have dimension $\dim U_f \cdot \dim W_f$. Since $\mathfrak{gl}_{U_f, W_f}$ injects into S_{U_f, W_f} under ι and does have dimension $\dim U_f \cdot \dim W_f$, we conclude that ι restricts to an isomorphism from $\mathfrak{gl}_{U_f, W_f}$ to S_{U_f, W_f}. Since any $\varphi \in S$ lies in some S_{U_f, W_f}, we have $\mathrm{im}\, \iota = S$. ∎

Corollary 3.2. If (U, W) is a finite-dimensional linear system, then $\mathfrak{gl}_{U,W} \cong \mathfrak{gl}^M_{U,W}$.

We identify $\mathfrak{gl}_{U_f,W_f}$ with S_{U_f,W_f}, and $\mathfrak{gl}_{U,W}$ with S inside $\mathfrak{gl}^M_{U,W}$ whenever appropriate.

Proposition 3.3. [PS13, Lemma 6.1] The Lie algebra $\mathfrak{gl}^M_{U,W}$ has a unique simple ideal $\mathfrak{sl}_{U,W}$.

Proposition 3.4. The Lie algebras $\mathfrak{gl}^M_{U_1,W_1}$ and $\mathfrak{gl}^M_{U_2,W_2}$ are isomorphic iff the linear systems (U_1, W_1) and (U_2, W_2) are isomorphic.

Proof. $\Longleftarrow$ This is obvious because the definition of $\mathfrak{gl}^M_{U,W}$ is intrinsic to the linear system (U, W). In the following we construct the isomorphism explicitly. If we have a linear system isomorphism $f\colon U_1 \to U_2$ and $g\colon W_1 \to W_2$, then we first show that $f^*_{|W_2} = g^{-1}$ and $(f^{-1})^*_{|W_1} = g$. Indeed

$$f^*(\langle \cdot, w_2\rangle) = \langle f(\cdot), w_2\rangle = \langle f^{-1}f(\cdot), g^{-1}(w_2)\rangle = \langle \cdot, g^{-1}(w_2)\rangle \quad \forall w_2 \in W_2,$$

$$(f^{-1})^*(\langle \cdot, w_1\rangle) = \langle f^{-1}(\cdot), w_1\rangle = \langle ff^{-1}(\cdot), g(w_1)\rangle = \langle \cdot, g(w_1)\rangle \quad \forall w_1 \in W_1.$$

Consider the map that sends $\varphi \in \mathfrak{gl}^M(U_1, W_1)$ to $f\varphi f^{-1}$. By the above

$$(f\varphi f^{-1})^*_{|W_2} = ((f^{-1})^*\varphi^* f^*)_{|W_2} = g\varphi^* g^{-1},$$

where the right hand side keeps W_2 stable. This map is a homomorphism of Lie algebras between $\mathfrak{gl}^M_{U_1,W_1}$ and $\mathfrak{gl}^M_{U_2,W_2}$, which has an inverse that sends $\psi \in \mathfrak{gl}^M_{U_2,W_2}$ to $f^{-1}\psi f$.

If we have a linear system isomorphism $f\colon U_1 \to W_2$ and $g\colon W_1 \to U_2$, then the map that sends φ to $f\varphi f^{-1}$ is an isomorphism of Lie algebras $\mathfrak{gl}^M_{U_1,W_1} \cong \mathfrak{gl}^M_{W_2,U_2}$ by the similar argument as above. But the latter is isomorphic to $\mathfrak{gl}^M_{U_2,W_2}$.

$\Longrightarrow$ Conversely if $\mathfrak{gl}^M_{U_1,W_1} \cong \mathfrak{gl}^M_{U_2,W_2}$, then $\mathfrak{sl}_{U_1,W_1} \cong \mathfrak{sl}_{U_2,W_2}$ because they are both the unique simple ideals of the respective Lie algebras. By Theorem 2.6, the linear systems (U_1, W_1) and (U_2, W_2) are isomorphic. □

Corollary 3.5. If (U, W) is countable, then $\mathfrak{gl}^M_{U,W} \cong \mathfrak{gl}^M_{V,V_*}$.

In addition, we observe the following about $\mathfrak{gl}^M_{U,W}$.

Proposition 3.6.

1. $\mathfrak{gl}^M_{U,W} \subset \mathfrak{gl}^M_{U,U^*} = \mathrm{End}(U)$.
2. $\mathfrak{gl}^M_{U,W}$ has one dimensional center $\mathbb{C}\mathrm{Id}$.
3. $\mathfrak{gl}_{U,W} \oplus \mathbb{C}\mathrm{Id}$ is an ideal of $\mathfrak{gl}^M_{U,W}$.

Proof.

1. Obvious.
2. Assume to the contrary that φ is in the center of $\mathfrak{gl}^M_{U,W}$ and $\varphi \notin \mathbb{C}\mathrm{Id}$. Then there exists $u \in U$ such that $\varphi(u) = u'$ and $u \neq \lambda u'$ for any $\lambda \in \mathbb{C}$, in particular $u \neq 0$. Extend the linearly independent set $\{u, u'\}$ to a basis of U, and define a linear operator ψ such that $\psi(u') = u$ and $\psi(v) = 0$ for any v in the basis that does not equal u'. Clearly $\psi \in \mathfrak{gl}_{U,W} \subset \mathfrak{gl}^M_{U,W}$, but
$$\varphi \circ \psi(u) = 0 \neq u = \psi \circ \varphi(u).$$
Contradiction.

3. $\mathbb{C}\mathrm{Id} \cap \mathfrak{gl}_{U,W} = 0$. The sum of two ideals is an ideal. □

Define $\mathfrak{gl}_\infty^M := \mathfrak{gl}_{V,V_*}^M$. Then $\mathfrak{gl}_\infty^M$ is isomorphic to the Lie algebra $\mathfrak{gl}_{U,W}^M$ for any countable dimensional linear system (U, W). In fact, $\mathfrak{gl}_\infty^M$ can be conveniently thought of as a certain Lie algebra of matrices.

Define $\mathrm{Mat}_{\mathbb{N}}^M$ to be the vector space consisting of matrices $(A_{ij})_{i,j\in\mathbb{N}}$ such that each row and each column has finitely many nonzero entries. Then $\mathrm{Mat}_{\mathbb{N}}^M$ is a Lie algebra and there is an isomorphism of Lie algebras $\mathfrak{gl}_\infty^M \cong \mathrm{Mat}_{\mathbb{N}}^M$.

We obtain the following commutative diagrams

$$\begin{array}{ccccc} \mathfrak{sl}_\infty & \hookrightarrow & \mathfrak{gl}_\infty & \hookrightarrow & \mathfrak{gl}_\infty^M \\ \updownarrow & & \updownarrow & & \updownarrow \\ \mathrm{Mat}_{\mathbb{N}}^0 & \hookrightarrow & \mathrm{Mat}_{\mathbb{N}} & \hookrightarrow & \mathrm{Mat}_{\mathbb{N}}^M. \end{array}$$

We know that V and V^* have dual $\mathfrak{gl}_\infty$-module structures and V_* is a submodule of V^*. Generalizing this, we give U and U^* dual $\mathfrak{gl}_{U,W}^M$-module structures, and let W be a submodule of U^*. By restriction, U and W become $\mathfrak{gl}_{U,W}$ and $\mathfrak{sl}_{U,W}$ modules as well. The actions are explicitly given by

$$\begin{aligned} &\varphi \cdot u = \varphi(u), \quad \varphi \cdot w = -\varphi^*(w) \quad \text{for } \varphi \in \mathfrak{gl}_{U,W}^M, u \in U, w \in W, \\ &u \otimes w \cdot x = \langle x, w\rangle u, \quad u \otimes w \cdot y = -\langle u, y\rangle w \quad \text{for } u \otimes w \in \mathfrak{gl}_{U,W}, x \in U, y \in W. \end{aligned}$$

Note that for any nonnegative integers p, q, the tensor product $U^{\otimes p} \otimes W^{\otimes q}$ also becomes a $\mathfrak{gl}_{U,W}^M$-module. Moreover, U and W are not isomorphic as $\mathfrak{gl}_{U,W}^M, \mathfrak{gl}_{U,W}$ or $\mathfrak{sl}_{U,W}$-modules, simply because $U \ncong W$ as $\mathfrak{sl}_{U_f,W_f}$-modules for finite-dimensional subsystems (U_f, W_f) with $\dim U_f = \dim W_f \geq 3$. Indeed, we know the following decomposition of $\mathfrak{sl}_{U_f,W_f}$-modules

$$U \cong U_f \oplus \left(\bigoplus_{n\in\mathbb{N}} \mathbb{C}\right) \quad \text{and} \quad W \cong W_f \oplus \left(\bigoplus_{n\in\mathbb{N}} \mathbb{C}\right),$$

but $W_f \cong U_f^* \ncong U_f$ as $\mathfrak{sl}_{U_f,W_f}$-modules if $\mathfrak{sl}_{U_f,W_f} \ncong \mathfrak{sl}(2)$.

4 The categories $\mathbb{T}_{\mathfrak{sl}_{U,W}}$, $\mathbb{T}_{\mathfrak{gl}_{U,W}^M}$ and the functor $\mathcal{F}^h$

In this section we present results about the categories $\mathbb{T}_{\mathfrak{sl}_{U,W}}$ and $\mathbb{T}_{\mathfrak{gl}_{U,W}^M}$ introduced in [PS13]. In particular, an object in $\mathbb{T}_{\mathfrak{sl}_{U,W}}$ or $\mathbb{T}_{\mathfrak{gl}_{U,W}^M}$ is isomorphic to a finite length submodule of a direct sum of finitely many copies of $T(U \oplus W)$. Moreover, the categories of tensor modules $\mathbb{T}_{\mathfrak{sl}_{U,W}}$ and $\mathbb{T}_{\mathfrak{gl}_{U,W}^M}$ are both equivalent to $\mathbb{T}_{\mathfrak{sl}_\infty}$ as monoidal categories.

Definition. Let M be an $\mathfrak{sl}_{U,W}$-module, we say M is $\mathfrak{sl}_{U,W}$-integrable iff for every $\varphi \in \mathfrak{sl}_{U,W}, m \in M$, we have $\dim\{m, \varphi \cdot m, \varphi^2 \cdot m, \cdots\} < \infty$.

A subalgebra $\mathfrak{k}$ of $\mathfrak{sl}_{U,W}$ has *finite co-rank* iff it contains $\mathfrak{sl}_{W_f^\perp, U_f^\perp}$ for some finite-dimensional subsystem (U_f, W_f). We say that a $\mathfrak{sl}_{U,W}$-module M satisfies the *large annihilator condition* if for every $m \in M$, its annihilator $\mathrm{Ann}(m) := \{g \in \mathfrak{sl}(U,W) \colon g \cdot m = 0\}$ is a finite co-rank subalgebra of $\mathfrak{sl}_{U,W}$.

Define $\mathbb{T}_{\mathfrak{sl}_{U,W}}$ to be the full subcategory of $\mathfrak{sl}_{U,W}$-mod where the objects are finite length $\mathfrak{sl}_{U,W}$-integrable modules that satisfy the large annihilator condition.

Despite the abstract definition, the category $\mathbb{T}_{\mathfrak{sl}_{U,W}}$ is nothing but the category of tensor modules.

Theorem 4.1. [PS13, Corollay 5.12] The following are equivalent:

1. M is an object of $\mathbb{T}_{\mathfrak{sl}_{U,W}}$.
2. M is isomorphic to a finite length submodule of a direct sum of finitely many copies of $T(U \oplus W)$;
3. M is isomorphic to a finite length subquotient of a direct sum of finitely many copies of $T(U \oplus W)$.

Theorem 4.2. [PS13, Theorem 5.5] For any linear system (U, W), there is an equivalence $\mathbb{T}_{\mathfrak{sl}_{U,W}} \leftrightsquigarrow \mathbb{T}_{\mathfrak{sl}_\infty}$ of monoidal tensor categories.

The category $\mathbb{T}_{\mathfrak{gl}^M_{U,W}}$ is defined analogously, but with slight differences. A subalgebra $\mathfrak{k}$ of $\mathfrak{gl}^M_{U,W}$ is called *finite co-rank* iff $\mathfrak{k} \supset \mathfrak{gl}^M_{W_f^\perp, U_f^\perp}$ for some finite-dimensional subsystem (U_f, W_f). A $\mathfrak{gl}^M_{U,W}$-module M is said to satisfy the *large annihilator condition* if $\mathrm{Ann}(m)$ is a finite co-rank subalgebra of $\mathfrak{gl}^M_{U,W}$ for any $m \in M$.

We define $\mathbb{T}_{\mathfrak{gl}^M_{U,W}}$ to be the full subcategory of $\mathfrak{gl}^M_{U,W}$-mod whose objects are finite length $\mathfrak{sl}_{U,W}$-integrable modules that satisfy the large annihilator condition.

Similar to the $\mathbb{T}_{\mathfrak{sl}_{U,W}}$ case, the category $\mathbb{T}_{\mathfrak{gl}^M_{U,W}}$ is nothing but the category of tensor modules.

Theorem 4.3. [PS13, Theorem 7.9 a)] The following are equivalent:

1. M is an object of $\mathbb{T}_{\mathfrak{gl}^M_{U,W}}$;
2. M is isomorphic to a finite length submodule of a direct sum of finitely many copies of $T(U \oplus W)$;
3. M is isomorphic to a finite length subquotient of a direct sum of finitely many copies of $T(U \oplus W)$.

Theorem 4.4. [PS13, Theorem 7.9 b)] For any object M in the category $\mathbb{T}_{\mathfrak{gl}^M_{U,W}}$, let $\mathrm{Res}(M)$ denote M regarded as a module of $\mathfrak{sl}_{U,W}$. Then Res: $\mathbb{T}_{\mathfrak{gl}^M_{U,W}} \rightsquigarrow \mathbb{T}_{\mathfrak{sl}_{U,W}}$ is a well defined functor that is fully faithful and essentially surjective. Moreover, $\mathrm{Res}(M \otimes N) = \mathrm{Res}(M) \otimes \mathrm{Res}(M)$. Thus $\mathbb{T}_{\mathfrak{gl}^M_{U,W}}$ and $\mathbb{T}_{\mathfrak{sl}_{U,W}}$ are equivalent as monoidal tensor categories.

We introduce the notion of *modules twisted by an automorphism*, or *twisted modules*. Let M be a $\mathfrak{g}$-module, and $h \in \mathrm{Aut}(\mathfrak{g})$ be an automorphism of $\mathfrak{g}$. Then the module twisted by h is the same as M as a vector space but with new action

$$\varphi \cdot_{new} w = h(\varphi) \cdot_{old} w \quad \forall \varphi \in \mathfrak{g}, \forall w \in M.$$

We denote this new $\mathfrak{g}$-module structure by M^h . In the language of representations, the pull-back of the representation $\rho\colon \mathfrak{g} \to \mathfrak{gl}(M)$ along $h\colon \mathfrak{g} \to \mathfrak{g}$ gives the representation $\rho \circ h$, which corresponds to the twisted module M^h.

Recall that the *socle* of a module M, denoted by $\operatorname{soc} M$, is the sum of simple submodules of M. If $f : M \to N$ is a $\mathfrak{g}$-module isomorphism, then clearly $f(\operatorname{soc} M) = \operatorname{soc} N$.

We observe the following about twisted modules.

Proposition 4.5. Let U, W be $\mathfrak{g}$-modules and $g, h \in \operatorname{Aut}(\mathfrak{g})$. Then the following holds:

1. $(W^h)^g = W^{g\circ h}$ as $\mathfrak{g}$-modules.
2. If $f : W \to U$ is a $\mathfrak{g}$-module homomorphism, then $f : W^h \to U^h$ is also a $\mathfrak{g}$-module homomorphism. In particular, if there is an isomorphism of $\mathfrak{g}$-modules $W \cong U$, then there is also an isomorphism of $\mathfrak{g}$-modules $W^h \cong U^h$.
3. Any $h \in \operatorname{Aut}(\mathfrak{g})$ induces a covariant functor $\mathcal{F}^h$ that sends W to W^h, and a morphism $f : W \to U$ to $f : W^h \to U^h$. The functor $\mathcal{F}^h$ has an inverse functor $\mathcal{F}^{h^{-1}}$, thus it is an automorphism of the category $\mathfrak{g}$-mod.
4. The functor $\mathcal{F}^h$ commutes with the contravariant functor $(\,\cdot\,)^*$, that is, $(U^*)^h = (U^*)^h$.
5. The functor $\mathcal{F}^h$ preserves the socle of a module, that is, $\operatorname{soc} W^h = (\operatorname{soc} W)^h$.

Proof.

1. The $\mathfrak{g}$-modules $(W^h)^g$ and $W^{g\circ h}$ have the same action

$$\varphi \cdot_{new} w = g(h(\varphi)) \cdot_{old} w \quad \forall \varphi \in \mathfrak{g}, w \in W.$$

2. Since $f\colon W \to U$ is a $\mathfrak{g}$-homomorphism, for every $\varphi \in \mathfrak{g}$ we have the following commutative diagram

$$\begin{array}{ccc} W & \xrightarrow{f} & U \\ {\scriptstyle\varphi}\downarrow & & \downarrow{\scriptstyle\varphi} \\ W & \xrightarrow{f} & U. \end{array}$$

   Then for every $\varphi \in \mathfrak{g}$ we also have the following commutative diagram

$$\begin{array}{ccc} W^h & \xrightarrow{f} & U^h \\ {\scriptstyle h(\varphi)}\downarrow & & \downarrow{\scriptstyle h(\varphi)} \\ W^h & \xrightarrow{f} & U^h. \end{array}$$

   Thus $f\colon W^h \to U^h$ is a $\mathfrak{g}$-module homomorphism.
3. It is clear from statement 2 that $\mathcal{F}^h$ takes a module W to W^h and morphisms $f\colon W \to U$ to the same morphism $f\colon W^h \to U^h$, and is a well-defined functor from $\mathfrak{g}$-mod to itself. The functors $\mathcal{F}^{h^{-1}}$ and $\mathcal{F}^h$ are mutually inverse by statement 1, thus $\mathcal{F}^h$ is an automorphism of the category $\mathfrak{g}$-mod.
4. Both actions are given by

$$(\varphi \cdot u^*)(u) = -u^*(h(\varphi) \cdot u) \quad \forall \varphi \in \mathfrak{g}, u \in U, u^* \in U^*.$$

5. Since $\mathcal{F}^h$ has an inverse $\mathcal{F}^{h^{-1}}$, it maps simple submodules to simple submodules. Also it is easy to see that for any submodules $W_1, W_2 \subset U$, we have $(W_1 + W_2)^h = W_1^h + W_2^h$. Therefore $\operatorname{soc} W^h = (\operatorname{soc} W)^h$. □

One important observation is that U^h and W^h are simple $\mathfrak{sl}_{U,W}$-modules for any $h \in \operatorname{Aut}(\mathfrak{sl}_{U,W})$. Further, we have $\operatorname{soc} (U^h)^* = W^h$ and $\operatorname{soc} (W^h)^* = U^h$.

Proposition 4.6. As $\mathfrak{sl}_{U,W}, \mathfrak{gl}_{U,W}$ and $\mathfrak{gl}_{U,W}^M$-modules,

1. U and W are simple.
2. $\operatorname{soc} U^* = W$ and $\operatorname{soc} W^* = U$.
3. $\operatorname{soc} (U^h)^* = W^h$ and $\operatorname{soc} (W^h)^* = U^h$ for any $h \in \operatorname{Aut}(\mathfrak{gl}_\infty^M)$.

Proof.

1. It suffices to check the simplicity of U and W as $\mathfrak{sl}_{U,W}$-modules. Let $X \subset U$ be a nontrivial submodule of U, then it intersects nontrivially with some finite-dimensional subspace $U_f \subset U$. Since U_f is a simple $\mathfrak{sl}_{U_f,W_f}$-module, we must have $X \supset U_f$. Therefore we must have $X \supset U_f'$ for any finite-dimensional subspace $U_f' \supset U_f$. We conclude that $X = U$, since U is the union of all its finite-dimensional subspaces. The same argument applies to W.
2. Let X be a nontrivial $\mathfrak{sl}_{U,W}$-submodule of U^*. It suffices to show that $X \supset W$. Since W is a simple module, all we need to show is that $X \cap W \neq 0$. Suppose the contrary, then take any nonzero element u^* in X, we must have $u^* \in U^* - W$. There exists some $u \in U$ such that $\langle u, u^* \rangle \neq 0$ by the non-degeneracy of the form $\langle \cdot, \cdot \rangle$. Pick some nonzero $w \in (\mathbb{C}u)^\perp \subset W$, then $u \otimes w \in \mathfrak{sl}_{U,W}$. However $u \otimes w \cdot u^* = \langle u, u^* \rangle w \in W \cap X$. Contradiction. The same argument can be applied to prove that $\operatorname{soc} W^* = U$.
3. By Proposition 4.5 statement 4 and 5,

$$\operatorname{soc} (U^h)^* = \operatorname{soc} (U^*)^h = (\operatorname{soc} U^*)^h = W^h,$$

$$\operatorname{soc} (W^h)^* = \operatorname{soc} (W^*)^h = (\operatorname{soc} W^*)^h = U^h.$$ □

5 Automorphisms of $\mathfrak{sl}_\infty$ and $\mathfrak{gl}_\infty^M$

In this section we investigate the automorphism groups of $\mathfrak{sl}_\infty$ and $\mathfrak{gl}_\infty^M$.

We say the linear system (U, W) is *self-dual* if there is an isomorphism of vector spaces $f\colon U \to W$ such that $\langle f^{-1}(w), f(u) \rangle = \langle u, w \rangle$ for every $u \in U, w \in W$. In this case, according to the proof of Proposition 3.4, we observe that

$$f^*_{|U} = f, \quad (f^{-1})^*_{|W} = f^{-1}.$$

By Proposition 3.4, there is an isomorphism of Lie algebras $\gamma\colon \mathfrak{gl}_{U,W}^M \to \mathfrak{gl}_{W,U}^M$ defined by $\gamma\colon \varphi \mapsto f\varphi f^{-1}$. The Lie algebra $\mathfrak{gl}_{W,U}^M$ is again isomorphic to $\mathfrak{gl}_{U,W}^M$ by the map $\sigma\colon \varphi \mapsto -\varphi^*_{|U}$.

The composition $\tau := \sigma\gamma$ is an automorphism of $\mathfrak{gl}^M_{U,W}$, which is explicitly

$$\tau\colon \varphi \mapsto -f^{-1}\varphi^* f \quad \forall \varphi \in \mathfrak{gl}^M_{U,W}.$$

From the definition of τ we see that τ is an involution. Moreover, for every $\varphi \in \mathfrak{gl}^M_{U,W}$ the following diagram commutes

$$\begin{array}{ccc} U & \xrightarrow{f} & W \\ {\scriptstyle \tau(\varphi)}\downarrow & & \downarrow{\scriptstyle -\varphi^*} \\ U & \xrightarrow{f} & W. \end{array}$$

Therefore we conclude that $f : U^\tau \to W$ is an isomorphism of $\mathfrak{gl}^M_{U,W}$-modules. Consequently $W^\tau \cong (U^\tau)^\tau = U$ as $\mathfrak{gl}^M_{U,W}$-modules. Since the Lie algebra $\mathfrak{gl}^M_{U,W}$ has a unique simple ideal $\mathfrak{sl}_{U,W}$, the automorphism τ restricts to an automorphism of $\mathfrak{sl}_{U,W}$.

If the linear system (U, W) has a pair of dual bases relative to $\langle\cdot,\cdot\rangle$, then clearly (U, W) is self-dual. The converse is false. Again, consider the linear system (X, X') which is defined at the end of section 2. Let $f : X \to X'$ be the isomorphism given by $f(w, w^*) = (w, w^*)$. Then $\langle f^{-1}(x), f(y)\rangle = \langle y, x\rangle$ for every $x \in X', y \in X$. Therefore the linear system (X, X') is self-dual, but it does not have a pair of dual bases.

In the countable dimensional case, the linear system (V, V_*) has a pair of dual bases. Let $\epsilon : V \to V_*$ be the isomorphism of vector spaces induced by $\epsilon(e_n) = e^n\ \forall e_n \in \mathcal{B}$. Then we observe that $\tau : \varphi \mapsto -\epsilon^{-1}\varphi^*\epsilon$ is the involution of the Lie algebra $\mathrm{Mat}^M_{\mathbb{N}}$ given by $A \mapsto -A^t$.

In the following, let $\mathfrak{g} = \mathfrak{sl}_\infty, \mathfrak{gl}^M_\infty$. Define $\tilde{G} := \{g \in \mathrm{Aut}(V) |\ g^*(V_*) = V_*\}$. Conjugation by an element in $\tilde{G}$ induces an automorphism of $\mathfrak{g}$. Therefore we have a group representation $\rho : \tilde{G} \to \mathrm{Aut}(\mathfrak{g})$ with kernel $\mathbb{C}^*$, where $\mathbb{C}^*$ denotes nonzero complex numbers. We set $\tilde{G}_0 := \mathrm{im}\ \rho$.

Theorem 5.1 (Beidar et.al, [BBCM02])**.** The subgroup $\tilde{G}_0$ has index 2 in $\mathrm{Aut}(\mathfrak{sl}_\infty)$, the quotient is represented by the involution τ. In fact, $\mathrm{Aut}(\mathfrak{sl}_\infty) = \tilde{G}_0 \rtimes \{\mathrm{Id}, \tau\}$.

Corollary 5.2. If $h \in \tilde{G}_0$, then $V^h \cong V$ and $V^{\tau\circ h} \cong V_*$ as $\mathfrak{sl}_\infty$-modules.

Proof. If $h \in \tilde{G}_0$, then $h(\varphi) = g^{-1}\varphi g$ for every $\varphi \in \mathfrak{sl}_\infty$ where $g \in \tilde{G}$. It follows that $g : V^h \to V$ is an automorphism of $\mathfrak{sl}_\infty$-modules. Therefore $V^{\tau\circ h} = (V^h)^\tau \cong V^\tau \cong V_*$ as $\mathfrak{sl}_\infty$-modules. □

Let M be a $\mathfrak{gl}^M_\infty$-module. We say that $\mathfrak{sl}_\infty$ *acts densely* on M, if for every $\varphi \in \mathfrak{gl}^M_\infty$ and any choice of finitely many vectors $r_1, \cdots, r_n \in M$, there exists $\psi \in \mathfrak{sl}_\infty$ such that $\psi \cdot r_k = \varphi \cdot r_k$ for $k = 1, \cdots, n$.

Lemma 5.3. [PS13, Lemma 8.2] Let L and L' be $\mathfrak{gl}^M_\infty$-modules which have finite length as $\mathfrak{sl}_\infty$-modules. Then

$$\mathrm{Hom}_{\mathfrak{sl}_\infty}(L, L') = \mathrm{Hom}_{\mathfrak{gl}^M_\infty}(L, L').$$

In particular, if L and L' are isomorphic as $\mathfrak{sl}_\infty$-modules, then they are isomorphic as $\mathfrak{gl}^M_\infty$-modules.

Proposition 5.4. [PS13, Proposition 8.1] Let M be a finite length $\mathfrak{gl}_\infty^M$-module that is $\mathfrak{sl}_\infty$-integrable. Then M is an object of $\mathbb{T}_{\mathfrak{gl}_\infty^M}$ iff $\mathfrak{sl}_\infty$ acts densely on it.

Corollary 5.5. Let $h \in \mathrm{Aut}(\mathfrak{gl}_\infty^M)$. Then there exists an isomorphism of $\mathfrak{gl}_\infty^M$-modules $V^h \cong V$ or $V^h \cong V_*$.

Proof. If $h \in \mathrm{Aut}(\mathfrak{gl}_\infty^M)$, then h restricts to an automorphism of $\mathfrak{sl}_\infty$. We show that $\mathfrak{sl}_\infty$ acts densely on V^h. For any $\varphi \in \mathfrak{gl}_\infty^M$ and $r_1, \cdots, r_n \in V$, we can find an element $\eta \in \mathfrak{sl}_\infty$ such that $\eta(r_k) = h(\varphi)(r_k)$ for $k = 1, \cdots, n$. Therefore $\psi := h^{-1}(\eta)$ is an element of $\mathfrak{sl}_\infty$ such that $\psi \cdot r_k = \varphi \cdot r_k$ for $k = 1, \cdots, n$. By Proposition 5.4, we conclude that $V^h \in \mathbb{T}_{\mathfrak{gl}_\infty^M}$. We know from Corollary 5.2 that either $V^h \cong V$ or $V^h \cong V_*$ as $\mathfrak{sl}_\infty$-modules, therefore we conclude that either $V^h \cong V$ or $V^h \cong V_*$ as $\mathfrak{gl}_\infty^M$-modules by Lemma 5.3. □

Theorem 5.6. Let $h \in \mathrm{Aut}(\mathfrak{gl}_\infty^M)$.

1. If there is an isomorphism $V^h \cong V$ of $\mathfrak{gl}_\infty^M$-modules, then $h \in \tilde{G}_0$.
2. If there is an isomorphism $V^h \cong V_*$ of $\mathfrak{gl}_\infty^M$-modules, then $\tau \circ h \in \tilde{G}_0$.

Proof.

1. If $f \colon V^h \to V$ is an $\mathfrak{gl}_\infty^M$-isomorphism, then $fh(\varphi) = \varphi f$ for every $\varphi \in \mathfrak{gl}_\infty^M$. Since f is an isomorphism, we conclude that $h(\varphi) = f^{-1}\varphi f$. What remains to be shown is that $f \in \tilde{G}$.

   Observe that the dual operator $f^* \colon V^* \to (V^h)^*$ is a $\mathfrak{gl}_\infty^M$-isomorphism. Since $\mathrm{soc}\, V^* = V_*$ and $\mathrm{soc}\, (V^h)^* = V_*^h$ by Proposition 4.6, we conclude that V_* is f^*-stable. Therefore, $f \in \tilde{G}$ and $h \in \tilde{G}_0$.
2. If $V^h \cong V_*$, then we have the following isomorphisms of $\mathfrak{gl}_\infty^M$-modules

$$V^{\tau \circ h} = (V^h)^\tau \cong V_*^\tau \cong (V^\tau)^\tau = V^{\tau^2} = V.$$

   By statement 1 we conclude that $\tau \circ h \in \tilde{G}_0$. □

Corollary 5.7.

1. $\mathrm{Aut}(\mathfrak{gl}_\infty^M) = \tilde{G}_0 \rtimes \{\mathrm{Id}, \tau\}$.
2. Every automorphism of $\mathfrak{sl}_\infty$ extends uniquely to an automorphism of $\mathfrak{gl}_\infty^M$.

Proof.

1. Follows directly from Corollary 5.5 and Theorem 5.6.
2. Follows directly from statement 1 and Theorem 5.1. □

In conclusion, we computed the automorphism group of $\mathfrak{gl}_\infty^M$ using knowledge about the categories $\mathbb{T}_{\mathfrak{sl}_\infty}$ and $\mathbb{T}_{\mathfrak{gl}_\infty^M}$. As a problem for the future, it would be interesting to determine the automorphism groups of $\mathfrak{sl}_{U,W}$, $\mathfrak{gl}_{U,W}$ and $\mathfrak{gl}_{U,W}^M$ for an arbitrary linear system (U, W).

References

[BBCM02] K.I. Beidar, M. Brësar, M.A. Chebotar, and W.S. Martindale, 3rd, On Herstein's Lie map conjectures, III. *J. Algebra* 249 (2002), 59-94.

[DCPS11] E. Dan-Cohen, I. Penkov, and V. Serganova, A Koszul category of representations of finitary Lie algebras. Preprint (2011), arXiv:1105.3407.

[Mac43] G. Mackey. On infinite dimensional linear spaces. *Trans. AMS* 57 (1945), 155-207.

[PS13] I. Penkov and V. Serganova. Tensor representations of Mackey Lie algebras and their dense subalgebras. *Developments and Retrospectives in Lie Theory: Algebraic Methods*, Springer Verlag (to appear) (2014).